

**A through D. Gill net CPUE (#/hr) by location, time period, and species in 2009.**

*A single net was set in each of the lake's five strata, on the bottom, in 4 to 5 meters of water, within one week of electrofishing, in spring and in fall.*

**a) Gill net CPUE by location, time period, and species.**

Year	Location	Season	Species	Mean CPUE
2009	Stratum 1	Spring	Smallmouth bass	0.488
		Spring	White perch	0.488
		Spring	Yellow perch	5.854
	Stratum 2	Spring	White perch	0.500
	Stratum 3	Spring	Alewife	1.500
		Spring	Gizzard shad	0.500
		Spring	White perch	1.000
		Spring	White sucker	0.500
		Spring	Yellow perch	0.500
	Stratum 4	Spring	Gizzard shad	0.500
		Spring	Walleye	2.500
		Spring	White perch	1.000
		Spring	White sucker	1.000
	Stratum 5	Spring	Gizzard shad	0.496
		Spring	Longnose gar	0.496
		Spring	Walleye	0.496
		Spring	White perch	1.488
		Spring	White sucker	0.496
	Stratum 1	Fall	Northern pike	0.500
	Stratum 2	Fall	White perch	0.500
	Stratum 3	Fall	Smallmouth bass	0.500
	Stratum 4	Fall	Largemouth bass	0.500
		Fall	Smallmouth bass	1.000
		Fall	White sucker	0.500
	Stratum 5	Fall	Channel catfish	0.500
Fall		Gizzard shad	3.000	
Fall		Yellow perch	0.500	

**A through D. Gill net CPUE (#/hr) by location, time period, and species in 2009 (continued).**

*A single net was set in each of the lake's five strata, on the bottom, in 4 to 5 meters of water, within one week of electrofishing, in spring and in fall.*

**b) Whole year gill net CPUE by location and species.**

Year	Location	Species	Mean CPUE
2009	Stratum 1	Northern pike	0.247
		Smallmouth bass	0.247
		White perch	0.247
		Yellow perch	2.963
	Stratum 2	White perch	0.500
	Stratum 3	Alewife	0.750
		Gizzard shad	0.250
		Smallmouth bass	0.250
		White perch	0.500
		White sucker	0.250
		Yellow perch	0.250
	Stratum 4	Gizzard shad	0.250
		Largemouth bass	0.250
		Smallmouth bass	0.500
		Walleye	1.250
		White perch	0.500
		White sucker	0.750
	Stratum 5	Channel catfish	0.249
		Gizzard shad	1.743
		Longnose gar	0.249
Walleye		0.249	
White perch		0.747	
White sucker		0.249	
Yellow perch		0.249	

**A through D. Gill net CPUE (#/hr) by location, time period, and species in 2009 (continued).**

*A single net was set in each of the lake's five strata, on the bottom, in 4 to 5 meters of water, within one week of electrofishing, in spring and in fall.*

**c) Gillnet CPUE by time period and species.**

Year	Season	Species	Mean CPUE	Relative Abundance
2009	Spring	Alewife	0.300	7.5%
		Gizzard shad	0.299	7.5%
		Longnose gar	0.099	2.5%
		Smallmouth bass	0.098	2.5%
		Walleye	0.599	15.0%
		White perch	0.895	22.5%
		White sucker	0.399	10.0%
		Yellow perch	1.271	32.5%
	Fall	Channel catfish	0.100	6.7%
		Gizzard shad	0.600	40.0%
		Largemouth bass	0.100	6.7%
		Northern pike	0.100	6.7%
		Smallmouth bass	0.300	20.0%
		White perch	0.100	6.7%
		White sucker	0.100	6.7%
Yellow perch	0.100	6.7%		

**A through D. Gill net CPUE (#/hr) by location, time period, and species in 2009 (continued).**

*A single net was set in each of the lake's five strata, on the bottom, in 4 to 5 meters of water, within one week of electrofishing, in spring and in fall.*

**d) Gillnet CPUE for entire year**

<b>Year</b>	<b>Species</b>	<b>Mean CPUE</b>	<b>Relative Abundance</b>
2009	Alewife	0.150	5.5%
	Channel catfish	0.050	1.8%
	Gizzard shad	0.449	16.4%
	Largemouth bass	0.050	1.8%
	Longnose gar	0.050	1.8%
	Northern pike	0.049	1.8%
	Smallmouth bass	0.199	7.3%
	Walleye	0.300	10.9%
	White perch	0.499	18.2%
	White sucker	0.250	9.1%
	Yellow perch	0.692	25.5%