

Library Reference 8.7.3.1.6

**Whole Lake Electrofishing number netted and number missed in 2009 for each sampling event (spring and fall) - 2009 entire year.**

*Note: Because of their large size carp are not boated, instead carp within netting distance are counted while still in the water therefore no carp within netting distance are ever "missed". Lepomis sp. include bluegill and pumpkinseed because these species cannot be distinguished while still in the water, at night when missed. Although most fish in the genus Micropterus sp. (black bass) are large enough to be readily distinguishable when missed, some are not, these are counted under the designation "Micropterus sp.". Bullhead that are missed are assumed to be brown bullhead. The "number netted" for shad and alewives (clupeids) differs from the "total number" depicted in Library Reference 8.7.3.1.5 (a-c) because the totals in those tables include both the netted and missed fish. For the clupeids in these tables, the sum of the "Number Missed" and "Number Netted" will equal the total number for those species in Library Reference 8.7.3.1.5 (a-c).*

a.) 2009 Entire year

<b>Species</b>	<b>Number Netted</b>	<b>Number Missed</b>
<b>Pumpkinseed</b>	1,606	977
<b>Yellow perch</b>	575	858
<b>Brown bullhead</b>	422	385
White perch	420	526
<b>Largemouth bass</b>	285	73
<b>Bluegill</b>	240	0
<b>Smallmouth bass</b>	163	0
White sucker	140	46
Carp	121	0
Gizzard shad	117	101
Golden shiner	113	5
<b>Rock bass</b>	87	0
Alewife	78	385
Bowfin	21	4
Shorthead redhorse	14	0
Freshwater drum	10	10
<b>Walleye</b>	10	4
Longnose gar	9	10
<b>Northern pike</b>	8	10
Yellow bullhead	7	0
<b>Channel catfish</b>	6	0
Black crappie	6	0
Banded killifish	4	0
Silver Redhorse Sucker	4	0
<i>Lepomis sp.</i>	2	1,510
Northern hog sucker	2	0
Chain pickerel	1	0
<b>Black bullhead</b>	1	0
Quillback	1	0
<i>Micropterus sp.</i>	0	72
Pike Family (Esocidae)	0	2

**Whole Lake Electrofishing number netted and number missed in 2009 for each sampling event (spring and fall) - 2009 entire year. (continued)**

*Note: Because of their large size carp are not boated, instead carp within netting distance are counted while still in the water therefore no carp within netting distance are ever "missed". Lepomis sp. include bluegill and pumpkinseed because these species cannot be distinguished while still in the water, at night when missed. Although most fish in the genus Micropterus sp. (black bass) are large enough to be readily distinguishable when missed, some are not, these are counted under the designation "Micropterus sp.". Bullhead that are missed are assumed to be brown bullhead. The "number netted" for shad and alewives (clupeids) differs from the "total number" depicted in Library Reference 8.7.3.1.5 (a-c) because the totals in those tables include both the netted and missed fish. For the clupeids in these tables, the sum of the "Number Missed" and "Number Netted" will equal the total number for those species in Library Reference 8.7.3.1.5 (a-c).*

b.) 2009 Spring

<b>Species</b>	<b>Number Netted</b>	<b>Number Missed</b>
<b>Pumpkinseed</b>	821	967
White perch	284	361
<b>Yellow perch</b>	266	397
<b>Brown bullhead</b>	263	290
<b>Largemouth bass</b>	120	64
<b>Smallmouth bass</b>	111	0
Carp	86	0
White sucker	76	32
Golden shiner	76	5
Alewife	75	385
<b>Bluegill</b>	70	0
Gizzard shad	21	10
<b>Rock bass</b>	18	0
Shorthead redhorse	11	0
<b>Walleye</b>	10	4
Longnose gar	8	9
Bowfin	7	3
<b>Northern pike</b>	4	3
Banded killifish	4	0
Freshwater drum	3	1
Northern hog sucker	2	0
Silver Redhorse Sucker	2	0
Yellow bullhead	2	0
<i>Lepomis sp.</i>	2	0
<b>Channel catfish</b>	1	0
Chain pickerel	1	0
Quillback	1	0
<b>Black bullhead</b>	1	0
Pike Family (Esocidae)	0	2

**Whole Lake Electrofishing number netted and number missed in 2009 for each sampling event (spring and fall) - 2009 entire year. (continued)**

*Note: Because of their large size carp are not boated, instead carp within netting distance are counted while still in the water therefore no carp within netting distance are ever "missed". Lepomis sp. include bluegill and pumpkinseed because these species cannot be distinguished while still in the water, at night when missed. Although most fish in the genus Micropterus sp. (black bass) are large enough to be readily distinguishable when missed, some are not, these are counted under the designation "Micropterus sp.". Bullhead that are missed are assumed to be brown bullhead. The "number netted" for shad and alewives (clupeids) differs from the "total number" depicted in Library Reference 8.7.3.1.5 (a-c) because the totals in those tables include both the netted and missed fish. For the clupeids in these tables, the sum of the "Number Missed" and "Number Netted" will equal the total number for those species in Library Reference 8.7.3.1.5 (a-c).*

c.) 2009 Fall

<b>Species</b>	<b>Number Netted</b>	<b>Number Missed</b>
<b>Pumpkinseed</b>	785	10
<b>Yellow perch</b>	309	461
<b>Bluegill</b>	170	0
<b>Largemouth bass</b>	165	9
<b>Brown bullhead</b>	159	95
White perch	136	165
Gizzard shad	96	91
<b>Rock bass</b>	69	0
White sucker	64	14
<b>Smallmouth bass</b>	52	0
Golden shiner	37	0
Carp	35	0
Bowfin	14	1
Freshwater drum	7	9
Black crappie	6	0
<b>Channel catfish</b>	5	0
Yellow bullhead	5	0
<b>Northern pike</b>	4	7
Alewife	3	0
Shorthead redhorse	3	0
Silver Redhorse Sucker	2	0
Longnose gar	1	1
<i>Lepomis sp.</i>	0	1,510
<i>Micropterus sp.</i>	0	72