

Library Reference 8.7.3.1.7

**A through O. Electrofishing CPUE (#/hr) and relative abundance in 2009 from each of five lake strata.**

*Note: CPUE for gamefish (bolded) is calculated from all transects within the Stratum. CPUE for non-gamefish are calculated from only the transects where all fish are collected. Because of the difficulty in netting clupeids (shad and alewives), the CPUE for these species is calculated from estimates of the number of fish.*

a.) Entire year 2009	Stratum 1				
	Species	Mean CPUE	SE	Relative abundance with clupeids	Relative abundance without clupeids
	Alewife	28.94	28.94	5.6%	-
	Black crappie	0.70	0.43	0.1%	0.2%
	<b>Bluegill</b>	29.95	8.41	5.8%	6.4%
	Bowfin	1.36	0.68	0.3%	0.3%
	<b>Brown bullhead</b>	42.27	7.11	8.2%	9.0%
	Carp	6.69	3.66	1.3%	1.4%
	Chain pickerel	0.35	0.35	0.1%	0.1%
	<b>Channel catfish</b>	0.35	0.35	0.1%	0.1%
	Gizzard shad	17.97	11.86	3.5%	-
	Golden shiner	27.67	10.55	5.4%	5.9%
	<b>Largemouth bass</b>	30.55	4.93	5.9%	6.5%
	Northern hog sucker	0.58	0.58	0.1%	0.1%
	<b>Northern pike</b>	1.13	0.73	0.2%	0.2%
	<b>Pumpkinseed</b>	171.58	33.44	33.2%	36.5%
	<b>Rock bass</b>	15.46	4.58	3.0%	3.3%
	Shorthead redhorse	0.58	0.58	0.1%	0.1%
	<b>Smallmouth bass</b>	10.85	3.18	2.1%	2.3%
	<b>Walleye</b>	0.35	0.35	0.1%	0.1%
	White perch	41.30	2.52	8.0%	8.8%
	White sucker	25.76	7.32	5.0%	5.5%
	Yellow bullhead	1.10	0.72	0.2%	0.2%
	<b>Yellow perch</b>	62.17	7.99	12.0%	13.2%

**A through O. Electrofishing CPUE (#/hr) and relative abundance in 2009 from each of five lake strata (continued).**

*Note: CPUE for gamefish (bolded) is calculated from all transects within the Stratum. CPUE for non-gamefish are calculated from only the transects where all fish are collected. Because of the difficulty in netting clupeids (shad and alewives), the CPUE for these species is calculated from estimates of the number of fish.*

b.) Spring 2009	Stratum 1				
	Species	Mean CPUE	SE	Relative abundance with clupeids	Relative abundance without clupeids
	Alewife	54.89	54.89	10.7%	-
	<b>Bluegill</b>	18.57	5.17	3.6%	4.1%
	<b>Brown bullhead</b>	56.83	12.05	11.0%	12.4%
	Carp	5.85	3.87	1.1%	1.3%
	Chain pickerel	0.66	0.66	0.1%	0.1%
	Gizzard shad	4.02	2.55	0.8%	-
	Golden shiner	40.24	14.47	7.8%	8.8%
	<b>Largemouth bass</b>	22.48	6.88	4.4%	4.9%
	Northern hog sucker	1.10	1.10	0.2%	0.2%
	<b>Northern pike</b>	1.64	1.01	0.3%	0.4%
	<b>Pumpkinseed</b>	151.99	35.81	29.5%	33.3%
	<b>Rock bass</b>	6.95	1.54	1.4%	1.5%
	Shorthead redhorse	1.10	1.10	0.2%	0.2%
	<b>Smallmouth bass</b>	18.68	6.23	3.6%	4.1%
	<b>Walleye</b>	0.66	0.66	0.1%	0.1%
	White perch	61.88	5.29	12.0%	13.6%
	White sucker	13.77	0.61	2.7%	3.0%
	<b>Yellow perch</b>	54.31	8.32	10.5%	11.9%

**A through O. Electrofishing CPUE (#/hr) and relative abundance in 2009 from each of five lake strata (continued).**

*Note: CPUE for gamefish (bolded) is calculated from all transects within the Stratum. CPUE for non-gamefish are calculated from only the transects where all fish are collected. Because of the difficulty in netting clupeids (shad and alewives), the CPUE for these species is calculated from estimates of the number of fish.*

c.) Fall 2009	Stratum 1				
	Species	Mean CPUE	SE	Relative abundance with clupeids	Relative abundance without clupeids
	Black crappie	1.40	0.86	0.3%	0.3%
	<b>Bluegill</b>	42.63	14.08	8.3%	8.8%
	Bowfin	2.73	1.37	0.5%	0.6%
	<b>Brown bullhead</b>	28.62	5.84	5.5%	5.9%
	Carp	7.63	3.53	1.5%	1.6%
	<b>Channel catfish</b>	0.73	0.73	0.1%	0.2%
	Gizzard shad	33.67	25.53	6.5%	-
	Golden shiner	13.88	7.84	2.7%	2.9%
	<b>Largemouth bass</b>	37.81	9.03	7.3%	7.8%
	<b>Northern pike</b>	0.67	0.67	0.1%	0.1%
	<b>Pumpkinseed</b>	191.68	35.99	37.1%	39.7%
	<b>Rock bass</b>	23.97	8.02	4.6%	5.0%
	<b>Smallmouth bass</b>	2.16	1.38	0.4%	0.5%
	White perch	19.07	10.56	3.7%	4.0%
	White sucker	37.41	14.04	7.2%	7.8%
	Yellow bullhead	2.16	1.38	0.4%	0.5%
	<b>Yellow perch</b>	70.13	9.49	13.6%	14.5%

**A through O. Electrofishing CPUE (#/hr) and relative abundance in 2009 from each of five lake strata (continued).**

*Note: CPUE for gamefish (bolded) is calculated from all transects within the Stratum. CPUE for non-gamefish are calculated from only the transects where all fish are collected. Because of the difficulty in netting clupeids (shad and alewives), the CPUE for these species is calculated from estimates of the number of fish.*

d.) Entire year 2009	Stratum 2				
	Species	Mean CPUE	SE	Relative abundance with clupeids	Relative abundance without clupeids
	Alewife	8.12	3.97	1.5%	-
	<b>Bluegill</b>	6.45	2.32	1.2%	2.8%
	Bowfin	1.39	0.71	0.3%	0.6%
	<b>Brown bullhead</b>	24.12	2.79	4.4%	10.6%
	<b>Brown trout</b>	0.40	0.40	0.1%	0.2%
	Carp	21.78	9.09	4.0%	9.6%
	Freshwater drum	2.26	1.27	0.4%	1.0%
	Gizzard shad	307.86	131.72	56.7%	-
	Golden shiner	2.21	2.21	0.4%	1.0%
	<b>Largemouth bass</b>	16.10	4.43	3.0%	7.1%
	<b>Pumpkinseed</b>	43.60	9.80	8.0%	19.2%
	<b>Rock bass</b>	4.05	1.48	0.8%	1.8%
	Shorthead redhorse	0.79	0.79	0.2%	0.4%
	<b>Smallmouth bass</b>	15.56	4.19	2.9%	6.9%
	<b>Walleye</b>	1.23	0.80	0.2%	0.5%
	White perch	21.15	7.42	3.9%	9.3%
	White sucker	45.00	10.04	8.3%	19.9%
	<b>Yellow perch</b>	20.58	6.67	3.8%	9.1%

**A through O. Electrofishing CPUE (#/hr) and relative abundance in 2009 from each of five lake strata (continued).**

*Note: CPUE for gamefish (bolded) is calculated from all transects within the Stratum. CPUE for non-gamefish are calculated from only the transects where all fish are collected. Because of the difficulty in netting clupeids (shad and alewives), the CPUE for these species is calculated from estimates of the number of fish.*

e.) Spring 2009	Stratum 2				
	Species	Mean CPUE	SE	Relative abundance with clupeids	Relative abundance without clupeids
	Alewife	231.58	62.00	29.8%	-
	Banded killifish	2.74	2.74	0.4%	0.5%
	<b>Bluegill</b>	8.10	2.41	1.0%	1.6%
	Bowfin	2.63	1.32	0.3%	0.5%
	<b>Brown bullhead</b>	60.58	5.05	7.8%	11.8%
	Carp	43.88	10.50	5.6%	8.6%
	Freshwater drum	2.77	1.39	0.4%	0.5%
	Gizzard shad	35.33	19.61	4.5%	-
	Golden shiner	6.94	3.70	0.9%	1.4%
	<b>Largemouth bass</b>	14.86	5.85	1.9%	2.9%
	Longnose gar	1.40	1.40	0.2%	0.3%
	<b>Northern pike</b>	0.64	0.64	0.1%	0.1%
	<b>Pumpkinseed</b>	97.47	26.21	12.5%	19.1%
	<b>Rock bass</b>	1.33	0.85	0.2%	0.3%
	Shorthead redhorse	9.51	3.79	1.2%	1.9%
	<b>Smallmouth bass</b>	14.28	2.86	1.8%	2.8%
	<b>Walleye</b>	1.37	1.37	0.2%	0.3%
	White perch	173.11	49.38	22.2%	33.8%
	White sucker	39.88	15.15	5.1%	7.8%
	<b>Yellow perch</b>	30.17	8.00	3.9%	5.9%

**A through O. Electrofishing CPUE (#/hr) and relative abundance in 2009 from each of five lake strata (continued).**

*Note: CPUE for gamefish (bolded) is calculated from all transects within the Stratum. CPUE for non-gamefish are calculated from only the transects where all fish are collected. Because of the difficulty in netting clupeids (shad and alewives), the CPUE for these species is calculated from estimates of the number of fish.*

f.) Fall 2009	Stratum 2				
	Species	Mean CPUE	SE	Relative abundance with clupeids	Relative abundance without clupeids
	Alewife	4.12	2.48	0.9%	-
	<b>Bluegill</b>	12.28	5.07	2.8%	4.6%
	<b>Brown bullhead</b>	24.99	5.53	5.7%	9.4%
	Carp	13.73	7.67	3.1%	5.2%
	Freshwater drum	4.93	2.47	1.1%	1.9%
	Gizzard shad	167.94	65.81	38.3%	-
	Golden shiner	2.40	2.40	0.6%	0.9%
	<b>Largemouth bass</b>	20.32	7.97	4.6%	7.6%
	Longnose gar	1.20	1.20	0.3%	0.5%
	<b>Northern pike</b>	0.60	0.60	0.1%	0.2%
	<b>Pumpkinseed</b>	50.96	23.54	11.6%	19.1%
	<b>Rock bass</b>	6.83	2.13	1.6%	2.6%
	Shorthead redhorse	1.43	1.43	0.3%	0.5%
	<b>Smallmouth bass</b>	9.89	6.03	2.3%	3.7%
	White perch	82.61	15.95	18.8%	31.0%
	White sucker	22.68	5.66	5.2%	8.5%
	<b>Yellow perch</b>	11.79	4.14	2.7%	4.4%

**A through O. Electrofishing CPUE (#/hr) and relative abundance in 2009 from each of five lake strata (continued).**

*Note: CPUE for gamefish (bolded) is calculated from all transects within the Stratum. CPUE for non-gamefish are calculated from only the transects where all fish are collected. Because of the difficulty in netting clupeids (shad and alewives), the CPUE for these species is calculated from estimates of the number of fish.*

g.) Entire year 2009	Stratum 3				
	Species	Mean CPUE	SE	Relative abundance with clupeids	Relative abundance without clupeids
	Alewife	160.43	87.91	31.5%	-
	<b>Bluegill</b>	9.09	2.71	1.8%	2.9%
	<b>Brown bullhead</b>	36.13	5.64	7.1%	11.5%
	Carp	10.61	4.93	2.1%	3.4%
	<b>Channel catfish</b>	1.10	0.66	0.2%	0.4%
	Freshwater drum	2.24	0.35	0.4%	0.7%
	Gizzard shad	34.50	3.42	6.8%	-
	Golden shiner	36.05	7.56	7.1%	11.5%
	<b>Largemouth bass</b>	15.52	4.07	3.0%	4.9%
	<i>Lepomis sp.</i>	0.95	0.95	0.2%	0.3%
	Northern hog sucker	0.95	0.95	0.2%	0.3%
	<b>Northern pike</b>	0.57	0.57	0.1%	0.2%
	<b>Pumpkinseed</b>	99.12	27.09	19.5%	31.5%
	<b>Rock bass</b>	3.21	2.15	0.6%	1.0%
	Shorthead redhorse	1.90	1.90	0.4%	0.6%
	<b>Smallmouth bass</b>	15.98	8.58	3.1%	5.1%
	<b>Walleye</b>	1.36	1.36	0.3%	0.4%
	White perch	33.02	20.07	6.5%	10.5%
	White sucker	27.61	8.65	5.4%	8.8%
	<b>Yellow perch</b>	19.36	4.60	3.8%	6.2%

**A through O. Electrofishing CPUE (#/hr) and relative abundance in 2009 from each of five lake strata (continued).**

*Note: CPUE for gamefish (bolded) is calculated from all transects within the Stratum. CPUE for non-gamefish are calculated from only the transects where all fish are collected. Because of the difficulty in netting clupeids (shad and alewives), the CPUE for these species is calculated from estimates of the number of fish.*

h.) Spring 2009	Stratum 3				
	Species	Mean CPUE	SE	Relative abundance with clupeids	Relative abundance without clupeids
	Alewife	336.67	200.46	47.6%	-
	<b>Bluegill</b>	5.48	1.42	0.8%	1.5%
	<b>Brown bullhead</b>	40.67	8.09	5.8%	11.0%
	Carp	6.15	6.15	0.9%	1.7%
	<b>Channel catfish</b>	0.95	0.95	0.1%	0.3%
	Gizzard shad	2.05	2.05	0.3%	-
	Golden shiner	57.65	28.46	8.2%	15.6%
	<b>Largemouth bass</b>	13.36	9.35	1.9%	3.6%
	<i>Lepomis sp.</i>	2.05	2.05	0.3%	0.6%
	Northern hog sucker	2.05	2.05	0.3%	0.6%
	<b>Northern pike</b>	1.07	1.07	0.2%	0.3%
	<b>Pumpkinseed</b>	108.59	47.59	15.3%	29.4%
	<b>Rock bass</b>	3.78	3.78	0.5%	1.0%
	Shorthead redhorse	4.10	4.10	0.6%	1.1%
	<b>Smallmouth bass</b>	15.91	8.96	2.3%	4.3%
	<b>Walleye</b>	2.84	2.84	0.4%	0.8%
	White perch	45.87	36.14	6.5%	12.4%
	White sucker	37.77	25.47	5.3%	10.2%
	<b>Yellow perch</b>	20.70	4.64	2.9%	5.6%



**A through O. Electrofishing CPUE (#/hr) and relative abundance in 2009 from each of five lake strata (continued).**

*Note: CPUE for gamefish (bolded) is calculated from all transects within the Stratum. CPUE for non-gamefish are calculated from only the transects where all fish are collected. Because of the difficulty in netting clupeids (shad and alewives), the CPUE for these species is calculated from estimates of the number of fish.*

i.) Fall 2009	Stratum 3				
	Species	Mean CPUE	SE	Relative abundance with clupeids	Relative abundance without clupeids
	<b>Bluegill</b>	12.55	4.60	3.7%	4.6%
	<b>Brown bullhead</b>	32.58	7.79	9.6%	12.0%
	Carp	16.61	16.61	4.9%	6.1%
	<b>Channel catfish</b>	1.38	1.38	0.4%	0.5%
	Freshwater drum	4.53	1.01	1.3%	1.7%
	Gizzard shad	66.73	0.27	19.7%	-
	Golden shiner	17.37	10.32	5.1%	6.4%
	<b>Largemouth bass</b>	18.52	3.48	5.5%	6.8%
	<b>Pumpkinseed</b>	93.91	28.63	27.7%	34.6%
	<b>Rock bass</b>	2.63	1.69	0.8%	1.0%
	<b>Smallmouth bass</b>	16.39	8.82	4.8%	6.0%
	White perch	22.41	5.80	6.6%	8.3%
	White sucker	15.11	9.57	4.5%	5.6%
	<b>Yellow perch</b>	17.81	5.96	5.3%	6.6%

**A through O. Electrofishing CPUE (#/hr) and relative abundance in 2009 from each of five lake strata (continued).**

*Note: CPUE for gamefish (bolded) is calculated from all transects within the Stratum. CPUE for non-gamefish are calculated from only the transects where all fish are collected. Because of the difficulty in netting clupeids (shad and alewives), the CPUE for these species is calculated from estimates of the number of fish.*

j.) Entire year 2009	Stratum 4				
	Species	Mean CPUE	SE	Relative abundance with clupeids	Relative abundance without clupeids
	Alewife	86.63	86.63	16.4%	-
	<b>Black bullhead</b>	0.43	0.43	0.1%	0.1%
	<b>Bluegill</b>	15.44	4.01	2.9%	3.5%
	<b>Brown bullhead</b>	23.57	8.56	4.5%	5.4%
	Carp	17.82	3.57	3.4%	4.1%
	<b>Channel catfish</b>	0.48	0.48	0.1%	0.1%
	Freshwater drum	1.19	1.19	0.2%	0.3%
	Gizzard shad	4.28	4.28	0.8%	-
	Golden shiner	6.42	6.42	1.2%	1.5%
	<b>Largemouth bass</b>	23.40	7.65	4.4%	5.4%
	Longnose gar	2.26	0.12	0.4%	0.5%
	<b>Northern pike</b>	0.44	0.44	0.1%	0.1%
	<b>Pumpkinseed</b>	149.40	28.15	28.3%	34.2%
	<b>Rock bass</b>	5.95	0.66	1.1%	1.4%
	Shorthead redhorse	2.14	2.14	0.4%	0.5%
	<b>Smallmouth bass</b>	14.84	3.98	2.8%	3.4%
	<b>Walleye</b>	1.79	0.88	0.3%	0.4%
	White perch	81.63	53.82	15.5%	18.7%
	White sucker	14.50	2.62	2.8%	3.3%
	Yellow bullhead	0.39	0.39	0.1%	0.1%
	<b>Yellow perch</b>	74.78	17.00	14.2%	17.1%

**A through O. Electrofishing CPUE (#/hr) and relative abundance in 2009 from each of five lake strata (continued).**

*Note: CPUE for gamefish (bolded) is calculated from all transects within the Stratum. CPUE for non-gamefish are calculated from only the transects where all fish are collected. Because of the difficulty in netting clupeids (shad and alewives), the CPUE for these species is calculated from estimates of the number of fish.*

k.) Spring 2009	Stratum 4				
	Species	Mean CPUE	SE	Relative abundance with clupeids	Relative abundance without clupeids
	Alewife	178.67	178.67	29.5%	-
	<b>Black bullhead</b>	0.88	0.88	0.2%	0.2%
	<b>Bluegill</b>	6.95	2.22	1.2%	1.6%
	<b>Brown bullhead</b>	25.16	8.59	4.2%	5.9%
	Carp	24.39	19.73	4.0%	5.8%
	Freshwater drum	2.33	2.33	0.4%	0.6%
	Gizzard shad	2.21	2.21	0.4%	-
	<b>Largemouth bass</b>	19.52	6.05	3.2%	4.6%
	Longnose gar	4.53	0.12	0.8%	1.1%
	<b>Pumpkinseed</b>	123.06	25.62	20.3%	29.0%
	<b>Rock bass</b>	1.81	1.11	0.3%	0.4%
	Shorthead redhorse	2.21	2.21	0.4%	0.5%
	<b>Smallmouth bass</b>	22.77	7.51	3.8%	5.4%
	<b>Walleye</b>	3.65	1.73	0.6%	0.9%
	White perch	101.35	61.65	16.8%	23.9%
	White sucker	20.34	1.71	3.4%	4.8%
	Yellow bullhead	0.78	0.78	0.1%	0.2%
	<b>Yellow perch</b>	64.59	15.10	10.7%	15.2%

**A through O. Electrofishing CPUE (#/hr) and relative abundance in 2009 from each of five lake strata (continued).**

*Note: CPUE for gamefish (bolded) is calculated from all transects within the Stratum. CPUE for non-gamefish are calculated from only the transects where all fish are collected. Because of the difficulty in netting clupeids (shad and alewives), the CPUE for these species is calculated from estimates of the number of fish.*

I.) Fall 2009	Stratum 4				
	Species	Mean CPUE	SE	Relative abundance with clupeids	Relative abundance without clupeids
	<b>Bluegill</b>	23.33	6.22	5.2%	5.2%
	<b>Brown bullhead</b>	22.34	9.28	4.9%	5.0%
	Carp	12.13	12.13	2.7%	2.7%
	<b>Channel catfish</b>	0.97	0.97	0.2%	0.2%
	Gizzard shad	6.23	6.23	1.4%	-
	Golden shiner	12.46	12.46	2.8%	2.8%
	<b>Largemouth bass</b>	27.12	9.47	6.0%	6.1%
	<b>Northern pike</b>	0.83	0.83	0.2%	0.2%
	<b>Pumpkinseed</b>	173.13	36.92	38.2%	38.8%
	<b>Rock bass</b>	9.96	1.49	2.2%	2.2%
	Shorthead redhorse	2.08	2.08	0.5%	0.5%
	<b>Smallmouth bass</b>	7.64	2.59	1.7%	1.7%
	White perch	61.67	45.06	13.6%	13.8%
	White sucker	8.65	3.80	1.9%	1.9%
	<b>Yellow perch</b>	84.33	23.27	18.6%	18.9%

**A through O. Electrofishing CPUE (#/hr) and relative abundance in 2009 from each of five lake strata (continued).**

*Note: CPUE for gamefish (bolded) is calculated from all transects within the Stratum. CPUE for non-gamefish are calculated from only the transects where all fish are collected. Because of the difficulty in netting clupeids (shad and alewives), the CPUE for these species is calculated from estimates of the number of fish.*

m.) Entire year 2009	Stratum 5				
	Species	Mean CPUE	SE	Relative abundance with clupeids	Relative abundance without clupeids
	Banded killifish	2.22	2.22	0.4%	0.4%
	Black crappie	2.05	1.19	0.4%	0.4%
	<b>Bluegill</b>	40.15	5.05	7.1%	7.1%
	<b>Brown bullhead</b>	34.97	4.35	6.2%	6.2%
	Carp	44.18	17.85	7.8%	7.8%
	<b>Channel catfish</b>	1.05	0.61	0.2%	0.2%
	Freshwater drum	1.01	1.01	0.2%	0.2%
	Gizzard shad	1.01	1.01	0.2%	-
	Golden shiner	22.31	2.06	4.0%	4.0%
	<b>Largemouth bass</b>	35.67	6.44	6.3%	6.3%
	<b>Northern pike</b>	0.52	0.52	0.1%	0.1%
	<b>Pumpkinseed</b>	204.07	17.67	36.2%	36.2%
	Quillback	1.01	1.01	0.2%	0.2%
	<b>Rock bass</b>	7.37	3.22	1.3%	1.3%
	Shorthead redhorse	1.01	1.01	0.2%	0.2%
	<b>Smallmouth bass</b>	15.22	3.34	2.7%	2.7%
	White perch	58.85	56.63	10.4%	10.5%
	White sucker	16.87	1.36	3.0%	3.0%
	Yellow bullhead	1.57	0.98	0.3%	0.3%
	<b>Yellow perch</b>	73.26	7.95	13.0%	13.0%

**A through O. Electrofishing CPUE (#/hr) and relative abundance in 2009 from each of five lake strata (continued).**

*Note: CPUE for gamefish (bolded) is calculated from all transects within the Stratum. CPUE for non-gamefish are calculated from only the transects where all fish are collected. Because of the difficulty in netting clupeids (shad and alewives), the CPUE for these species is calculated from estimates of the number of fish.*

n.) Spring 2009	Stratum 5				
	Species	Mean CPUE	SE	Relative abundance with clupeids	Relative abundance without clupeids
	Banded killifish	4.27	4.27	0.7%	0.7%
	<b>Bluegill</b>	21.07	7.69	3.5%	3.5%
	<b>Brown bullhead</b>	36.69	8.18	6.0%	6.0%
	Carp	71.57	39.46	11.8%	11.8%
	Golden shiner	20.84	4.78	3.4%	3.4%
	<b>Largemouth bass</b>	34.80	14.76	5.7%	5.7%
	<b>Pumpkinseed</b>	235.47	39.34	38.7%	38.7%
	Quillback	2.01	2.01	0.3%	0.3%
	<b>Rock bass</b>	1.07	1.07	0.2%	0.2%
	<b>Smallmouth bass</b>	22.90	5.29	3.8%	3.8%
	White perch	74.25	74.25	12.2%	12.2%
	White sucker	20.45	7.64	3.4%	3.4%
	Yellow bullhead	1.07	1.07	0.2%	0.2%
	<b>Yellow perch</b>	61.80	10.28	10.2%	10.2%

**A through O. Electrofishing CPUE (#/hr) and relative abundance in 2009 from each of five lake strata (continued).**

*Note: CPUE for gamefish (bolded) is calculated from all transects within the Stratum. CPUE for non-gamefish are calculated from only the transects where all fish are collected. Because of the difficulty in netting clupeids (shad and alewives), the CPUE for these species is calculated from estimates of the number of fish.*

o.) Fall 2009	Stratum 5				
	Species	Mean CPUE	SE	Relative abundance with clupeids	Relative abundance without clupeids
	Black crappie	4.11	2.37	0.8%	0.8%
	<b>Bluegill</b>	59.23	12.44	11.5%	11.5%
	<b>Brown bullhead</b>	33.07	5.91	6.4%	6.4%
	Carp	14.83	5.62	2.9%	2.9%
	<b>Channel catfish</b>	2.12	1.22	0.4%	0.4%
	Freshwater drum	2.05	2.05	0.4%	0.4%
	Gizzard shad	2.05	2.05	0.4%	-
	Golden shiner	23.78	0.76	4.6%	4.6%
	<b>Largemouth bass</b>	36.15	5.01	7.0%	7.0%
	<b>Northern pike</b>	1.03	1.03	0.2%	0.2%
	<b>Pumpkinseed</b>	170.13	25.92	33.0%	33.2%
	<b>Rock bass</b>	13.67	6.76	2.7%	2.7%
	Shorthead redhorse	2.05	2.05	0.4%	0.4%
	<b>Smallmouth bass</b>	7.44	2.64	1.4%	1.5%
	White perch	43.21	38.61	8.4%	8.4%
	White sucker	13.30	5.12	2.6%	2.6%
	Yellow bullhead	2.05	2.05	0.4%	0.4%
	<b>Yellow perch</b>	85.06	9.13	16.5%	16.6%