

Library Reference 8.7.3.2.7

**Relative contribution of each Pollution Tolerance category in 2009 for each electrofishing event (spring and fall) and both events combined.**

*Refer to AMP 2009 Library Reference 8.7.3.2.1 for a list of species within each guild.*

Area	Time	Tolerant	Moderately Tolerant	Moderate	Moderately Intolerant	Intolerant
Whole Lake	Spring	42%	27%	27%	4%	0%
	Fall	52%	24%	24%	0%	0%
	Whole Year	44%	26%	26%	4%	0%
Stratum 1	Spring	41%	29%	24%	6%	0%
	Fall	59%	24%	18%	0%	0%
	Whole Year	48%	29%	19%	5%	0%
Stratum 2	Spring	45%	25%	30%	0%	0%
	Fall	47%	24%	29%	0%	0%
	Whole Year	45%	25%	30%	0%	0%
Stratum 3	Spring	42%	32%	21%	5%	0%
	Fall	50%	29%	21%	0%	0%
	Whole Year	40%	30%	25%	5%	0%
Stratum 4	Spring	44%	33%	22%	0%	0%
	Fall	47%	33%	20%	0%	0%
	Whole Year	43%	33%	24%	0%	0%
Stratum 5	Spring	57%	14%	29%	0%	0%
	Fall	50%	28%	22%	0%	0%
	Whole Year	45%	25%	30%	0%	0%

Percent of species in each tolerance category, based on presence of species.

Tolerance designations based on Whittier and Hughes (1998)